


Fy

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07D 401/12, 231/14, 403/12, 405/12, 473/34, 417/12, A61K 31/415, 31/4155		A1	(11) International Publication Number: WO 00/69849
			(43) International Publication Date: 23 November 2000 (23.11.00)
(21) International Application Number: PCT/US00/11903 (22) International Filing Date: 2 May 2000 (02.05.00) (30) Priority Data: 60/133,842 12 May 1999 (12.05.99) US (71) Applicant: ORTHO-MCNEIL PHARMACEUTICAL, INC. [US/US]; 920 U.S. Route 202, Raritan, NJ 08869 (US). (72) Inventors: KORDIK, Cheryl, P.; 102 Summer Ridge Drive, Lansdale, PA 19446 (US). LOVENBERG, Timothy, W.; 13252 Courtland Terrace, SanDiego, CA 92130 (US). REITZ, Allen, B.; 109 Greenbrier Road, Lansdale, PA 19446 (US). (74) Agent: APPOLLINA, Mary, A.; Johnson & Johnson, One Johnson & Johnson Plaza, New Brunswick, NJ 08933 (US).		(81) Designated States: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(54) Title: PYRAZOLE CARBOXAMIDES USEFUL FOR THE TREATMENT OF OBESITY AND OTHER DISORDERS			
<div style="display: flex; align-items: center; justify-content: center;">  <div style="margin-left: 20px;">(I)</div> </div>			
(57) Abstract Pyrazole carboxamide derivatives of formula (I) which are ligands for the neuropeptide Y, subtype 5 receptor, and pharmaceutical compositions containing a pyrazole carboxamide derivative as the active ingredient are described. The pyrazole carboxamides are useful in the treatment of disorders and diseases associated with the NPY receptor subtype Y5.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Larvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

TITLE OF THE INVENTION

PYRAZOLE CARBOXAMIDES USEFUL FOR THE TREATMENT OF
OBESITY AND OTHER DISORDERS

5

CROSS-REFERENCE TO RELATED APPLICATION

This application claims priority from United States provisional application
Serial No. 60/133,842, filed May 12, 1999, the contents of which are hereby
10 incorporated by reference.

FIELD OF THE INVENTION

This invention relates to a series of pyrazole carboxamide derivatives,
pharmaceutical compositions containing them and their preparation use in the
15 treatment of central nervous system disorders and affective conditions. More
particularly, the compounds of the invention are ligands for the neuropeptide Y
Y5 (NPY5) receptor, a receptor which is associated with a number of central
nervous system disorders and affective conditions.

20 BACKGROUND OF THE INVENTION

Regulation and function of the mammalian central nervous system is
governed by a series of interdependent receptors, neurons, neurotransmitters,
and proteins. The neurons play a vital role in this system, for when externally
or internally stimulated, they react by releasing neurotransmitters that bind to
25 specific proteins. Common examples of endogenous small molecule
neurotransmitters such as acetylcholine, adrenaline, norepinephrine,
dopamine, serotonin, glutamate, and gamma-aminobutyric acid are well
known, as are the specific receptors that recognize these compounds as
ligands ("The Biochemical Basis of Neuropharmacology", Sixth Edition,
30 Cooper, J. R.; Bloom, F. E.; Roth, R. H. Eds., Oxford University Press, New
York, NY 1991).

In addition to the endogenous small molecule neurotransmitters, there is increasing evidence that neuropeptides play an integral role in neuronal operations. Neuropeptides are now believed to be co-localized with perhaps more than one-half of the 100 billion neurons of the human central nervous system. In addition to humans, neuropeptides have been discovered in a number of animal species. In some instances the composition of these peptides is remarkably homogenous among species. This finding suggests that the function of neuropeptides is vital and has been impervious to evolutionary changes. Furthermore, neuropeptides, unlike small molecule neurotransmitters, are typically synthesized by the neuronal ribosome. In some cases, the active neuropeptides are produced as part of a larger protein which is enzymatically processed to yield the active substance. Based upon these differences, compared to small molecule neurotransmitters, neuropeptide-based strategies may offer novel therapies for CNS diseases and disorders. Specifically, agents that affect the binding of neuropeptides to their respective receptors or ameliorate responses that are mediated by neuropeptides are potential therapies for diseases associated with neuropeptides.

There are a number of afflictions that are associated with the complex interdependent system of receptors and ligands within the central nervous system; these include neurodegenerative diseases, affective disorders such as anxiety, depression, pain and schizophrenia, and affective conditions that include a metabolic component, namely obesity. Such conditions, disorders and diseases have been treated with small molecules and peptides which modulate neuronal responses to endogenous neurotransmitters.

One example of the class of neuropeptides is neuropeptide Y (NPY). NPY was first isolated from porcine brain (Tatemoto, K. et al. *Nature* 1982, 296, 659) and was shown to be structurally similar to other members of the pancreatic polypeptide (PP) family such as peptide YY, which is primarily synthesized by endocrine cells in the gut, and pancreatic polypeptide, which is

synthesized by the pancreas. Neuropeptide Y is a single peptide protein that consists of thirty-six amino acids containing an amidated C-terminus. Like other members of the pancreatic polypeptide family, NPY has a distinctive conformation that consists of an N-terminal polyproline helical region and an
5 amphiphilic α -helix joined by a characteristic PP-fold (Vladimir, S. et. Al. *Biochemistry* **1990**, 20, 4509). Furthermore, NPY sequences from a number of animal species have been elucidated and all show a high degree of amino acid homology to the human protein (>94% in rat, dog, rabbit, pig, cow, sheep) (see Larhammar, D. in "The Biology of Neuropeptide Y and Related Peptides",
10 Colmers, W. F. and Wahlestedt, C. Eds., Humana Press, Totowa, NJ 1993).

Endogenous receptor proteins that bind NPY and related peptides as ligands have been identified and distinguished, and several such proteins have been cloned and expressed. Six different receptor subtypes [Y1, Y2, Y3, Y4(PP), Y5, Y6 (formerly designated as a Y5 receptor)] are recognized today
15 based upon binding profile, pharmacology and / or composition if identity is known (Wahlestedt, C. et. al. *Ann. NY Acad. Sci.* **1990**, 611, 7; Larhammar, D. et. al. *J. Biol. Chem.* **1992**, 267, 10935; Wahlestedt, C. et. al. *Regul. Pept.* **1986**, 13, 307; Fuhlendorff, J. U. et. al. *Proc. Natl. Acad. Sci. USA* **1990**, 87,
20 182; Grundemar, L. et. al. *J. Pharmacol. Exp. Ther.* **1991**, 258, 633; Laburthe, M. et. al. *Endocrinology* **1986**, 118, 1910; Castan, I. et. al. *Endocrinology* **1992**, 131, 1970; Gerald, C. et. al. *Nature* **1996**, 382, 168; Weinberg, D. H. et. al. *Journal of Biological Chemistry* **1996**, 271, 16435; Gehlert, D. et. al. *Current Pharmaceutical Design* **1995**, 1, 295; Lundberg, J. M. et. al. *Trends in*
25 *Pharmaceutical Sciences* **1996**, 17, 301). Most and perhaps all NPY receptor proteins belong to the family of so-called G-protein coupled receptors (GPCRs). The neuropeptide Y5 receptor, a putative GPCR, is negatively coupled to cellular cyclic adenosine monophosphate (cAMP) levels via the action of adenylate cyclase (Gerald, C. et. al. *Nature* **1996**, 382, 168; Gerald,
30 C. et. al. PCT WO 96/16542). For example, NPY inhibits forskolin-stimulated cAMP production / levels in a neuroblastoma cell line. A Y5 ligand that mimics NPY in this fashion is an agonist whereas one that competitively reverses the

NPY inhibition of forskolin-stimulated cAMP production is an antagonist.

Neuropeptide Y itself is the archetypal substrate for the NPY receptors and its binding can elicit a variety of pharmacological and biological effects *in vitro* and *in vivo*. When administered to the brain of live animals (intracerebroventricularly (icv) or into the amygdala), NPY produces anxiolytic effects in established animal models of anxiety such as the elevated plus-maze, Vogel punished drinking and Geller-Seifter's bar-pressing conflict paradigms (Heilig, M. et. al. *Psychopharmacology* 1989, 98, 524; Heilig, M. et. al. *Reg. Peptides* 1992, 41, 61; Heilig, M. et. al. *Neuropsychopharmacology* 1993, 8, 357). Thus compounds that mimic NPY are postulated to be useful for the treatment of anxiolytic disorders.

The immunoreactivity of neuropeptide Y is notably decreased in the cerebrospinal fluid of patients with major depression and those of suicide victims (Widdowson, P. S. et. al. *Journal of Neurochemistry* 1992, 59, 73), and rats treated with tricyclic antidepressants display significant increases of NPY relative to a control group (Heilig, M. et. al. *European Journal of Pharmacology* 1988, 147, 465). These findings suggest that an inadequate NPY response may play a role in some depressive illnesses, and that compounds that regulate the NPY-ergic system may be useful for the treatment of depression.

Neuropeptide Y improves memory and performance scores in animal models of learning (Flood, J. F. et. al. *Brain Research* 1987, 421, 280) and therefore may serve as a cognition enhancer for the treatment of neurodegenerative diseases such as Alzheimer's Disease (AD) as well as AIDS-related and senile dementia.

Elevated plasma levels of NPY are present in animals and humans experiencing episodes of high sympathetic nerve activity such as surgery, newborn delivery and hemorrhage (Morris, M. J. et. al. *Journal of Autonomic Nervous System* 1986, 17, 143). Thus chemical substances that alter the

NPY-ergic system may be useful for alleviating the condition of stress.

Neuropeptide Y also mediates endocrine functions such as the release of luteinizing hormone (LH) in rodents (Kalra, S. P. et. al. *Frontiers in Neuroendocrinology* **1992**, 13, 1). Since LH is vital for mammalian ovulation, a compound that mimics the action of NPY could be useful for the treatment of infertility, particularly in women with so-called luteal phase defects.

Neuropeptide Y is a powerful stimulant of food intake; as little as one-billionth of a gram, when injected directly into the CNS, causes satiated rats to overeat (Clark, J. T. et. al. *Endocrinology* **1984**, 115, 427; Levine, A. S. et. al. *Peptides* **1984**, 5, 1025; Stanley, B. G. et. al. *Life Sci.* **1984**, 35, 2635; Stanley, B. G. et. al. *Proc. Nat. Acad. Sci. USA* **1985**, 82, 3940). Thus NPY is orexigenic in rodents but not anxiogenic when given intracerebroventricularly and so antagonism of neuropeptide receptors may be useful for the treatment of eating disorders such as obesity, binge eating, anorexia nervosa and bulimia nervosa.

In recent years, a variety of potent, structurally distinct small molecule Y1 antagonists has been discovered and developed (Hipskind, P. A. et. al. *Annu. Rep. Med. Chem.* **1996**, 31, 1-10; Rudolf, K. et. al. *Eur. J. Pharmacol.* **1994**, 271, R11; Serradeil-Le Gal, C. et. al. *FEBS Lett.* **1995**, 362, 192; Wright, J. et. al. *Bioorg. Med. Chem. Lett.* **1996**, 6, 1809; Poindexter, G. S. et. al. United States Patent 5,668,151; Peterson, J. M. et. al. WO9614307 (1996)). However, despite claims of activity in rodent models of feeding, it is unclear if inhibition of a feeding response can be attributed to antagonism of the Y1 receptor.


Several landmark studies strongly suggest that an "atypical Y1" receptor and / or the Y5 receptor, rather than the classic Y1 receptor, is responsible for invoking NPY-stimulated food consumption in animals. It has been shown that the NPY fragment NPY2-36 is a potent inducer of feeding despite poor binding

at the classic Y1 receptor (Stanley, B. G. et. al. *Peptides* 1992, 13, 581). Conversely, a potent and selective Y1 agonist has been reported to be inactive at stimulating feeding in animals (Kirby, D. A. et. al. *J. Med. Chem.* 1995, 38, 4579). More pertinent to the invention described herein, [D-Trp³²]NPY, a
5 selective Y5 receptor activator has been reported to stimulate food intake when injected into the hypothalamus of rats (Gerald, C. et. al. *Nature* 1996, 382, 168). Since [D-Trp³²]NPY appears to be a full agonist of the Y5 receptor with no appreciable Y1 activity, the Y5 receptor is hypothesized to be responsible for the feeding response. Accordingly compounds that antagonize
10 the Y5 receptor should be effective in inhibiting food intake, particularly that stimulated by NPY.

Also pertinent to the invention described herein, are aminopyrazoles that act as Y5 antagonists. In PCT WO 98/27063 and PCT WO 98/25908,
15 certain aminopyrazoles are described as Y5 antagonists. In PCT WO 98/25907, (carbonylamino)pyrazole derivatives are likewise claimed as Y5 antagonists.

20 SUMMARY OF THE INVENTION

The present invention is directed to compounds of the formula (I):



(I)

wherein

25

R¹ is selected from the group consisting of C₂-C₈ alkyl, aryl, substituted

aryl, arC₁-C₈ alkyl, substituted arC₁-C₈ alkyl, heteroaryl, substituted heteroaryl, C₃-C₈ cycloalkyl, heteroC₃-C₈cycloalkyl, fluorinated C₁-C₈ alkyl, cyanoC₁-C₈alkyl and hydroxyC₁-C₈alkyl; wherein the aryl, aralkyl or heteroaryl group is substituted with one or more substituents independently selected from

5 halogen, C₁-C₈ alkyl, C₁-C₈ alkoxy, fluorinated C₁-C₈ alkyl, fluorinated C₁-C₈ alkoxy, nitro, amino, amido, N-C₁-C₈alkylamido, N,N-di(C₁-C₈alkyl)amido, C₁-C₈ alkylsulfonyl, sulfonamido, N-C₁-C₈alkylsulfonamido, N,N-di(C₁-C₈alkyl)sulfonamido, C₁-C₈alkylsulfonylamino, or C₁-C₈alkylcarbonylamino;

10

R² is selected from the group consisting of di(C₁-C₆ alkyl)amino-C₁-C₆ alkyl, unsubstituted or substituted heteroarylC₀-C₈ alkyl, unsubstituted or substituted heterocycloalkylC₀-C₆ alkyl, unsubstituted or substituted aryl, unsubstituted or substituted arC₁-C₆ alkyl, and unsubstituted or substituted

15 C₃-C₈ cycloalkylC₁-C₆ alkyl where the substituent is arylsulfonamidoC₁-C₆ alkyl;

wherein

- (a) the substituents on the aryl group are one or more substituents independently selected from the group consisting of
- 20 halogen, nitro, amino, substituted amino where the substituents on the amino are one or two of C₁-C₆ alkyl or phenyl; fluorinated C₁-C₆ alkyl, fluorinated C₁-C₆ alkoxy, C₁-C₆ alkyl, substituted C₁-C₆ alkyl where the substituent on the alkyl is N-(C₁-C₆ alkyl)-N-(C₃-C₈ cycloalkyl)amino; C₁-C₈ alkoxy, arC₁-C₈ alkyl, arC₁-C₈ alkoxy,
- 25 arylcarbonyl, phenyl, aminophenyl, C₁-C₆ alkylthio, substituted arylsulfonamido where the substituent on the arylsulfonamido is C₁-C₆ alkyl or C₁-C₆ alkoxy; heteroaryl, substituted heteroaryl where the substituent on the heteroaryl is C₁-C₆ alkyl, heterocycloalkyl and oxo;
- 30 (b) the substituents on the aralkyl group are one or more substituents independently selected from halogen or fluorinated C₁-C₄ alkyl;

(c) the substituents on the heterocycloalkyl group are one or more substituents independently selected from C₁-C₆ alkyl or arC₁-C₈ alkyl;

5

(d) the substituents on the heteroaryl group are one or more substituents independently selected from oxo, C₁-C₆ alkyl or halogen;

10 R³ is selected from the group consisting of hydrogen, halogen, C₁-C₈ alkyl, arC₁-C₈ alkyl, heteroarylC₁-C₈ alkyl, C₁-C₈ alkoxy, fluorinated C₁-C₈ alkyl, fluorinated C₁-C₈ alkoxy, aminoC₁-C₈ alkyl and C₁-C₈ alkylaminoC₁-C₈ alkyl; preferably, R³ is hydrogen;

15 R⁴ is selected from the group consisting of halogen, C₁-C₈ alkyl, arC₁-C₈ alkyl, heteroarylC₁-C₈ alkyl, fluorinated C₁-C₈ alkyl, aminoC₁-C₈ alkyl and C₁-C₈ alkylaminoC₁-C₈ alkyl; preferably, R⁴ is C₁-C₄ alkyl;

R⁵ is selected from the group consisting of hydrogen and C₁-C₈ alkyl;
20 preferably, R⁵ is hydrogen or C₁-C₄ alkyl

provided that when R¹ is phenyl or benzyl, and R³ is hydrogen, and R⁴ is methyl, and R⁵ is hydrogen, then R² is selected from di(C₁-C₆ alkyl)aminoC₁-C₆ alkyl, unsubstituted or substituted heteroarylC₀-C₆ alkyl, unsubstituted or substituted heterocycloalkylC₀-C₆ alkyl, substituted arC₁-C₆ alkyl, unsubstituted or substituted C₃-C₈ cycloalkylC₁-C₆ alkyl where the substituent is arylsulfonamidoC₁-C₆ alkyl, and substituted aryl wherein the substituents on the aryl group are one or more substituents independently selected from the group consisting of halogen, nitro, amino, substituted amino where the substituents on the amino are one or two of C₁-C₆ alkyl or phenyl; fluorinated C₁-C₆ alkyl, fluorinated C₁-C₆ alkoxy, C₁-C₆ alkyl (provided that R² is not di(C₁-C₄ alkyl)phenyl), substituted C₁-C₆ alkyl where the substituent on the alkyl is N-

30

(C₁-C₆ alkyl)-N-(C₃-C₈ cycloalkyl)amino; C₁-C₈ alkoxy, arC₁-C₈ alkyl, arC₁-C₈ alkoxy, phenyl, aminophenyl, C₁-C₆ alkylthio, substituted arylsulfonamido where the substituent on the arylsulfonamido is C₁-C₆ alkyl or C₁-C₆ alkoxy; heteroaryl, substituted heteroaryl where the substituent on the heteroaryl is C₁-C₆ alkyl; heterocycloalkyl and oxo;

and pharmaceutically acceptable salts thereof.

In a one embodiment of the invention are compounds of the formula (I) wherein

10 R¹ is selected from the group consisting of phenyl, 3-tolyl, 3-trifluoromethylphenyl, 3,5-dichlorophenyl, 3-chlorophenyl, and 2-chloro-5-trifluoromethylphenyl;

R² is selected from the group consisting of 4-fluorophenyl, 3,5-dichlorophenyl, 2,3-dichlorophenyl, 2-(1-pyrrolyl) phenyl, 2-(aminophenyl)phenyl, 3-trifluoromethoxyphenyl, 2-methylthio, 3-methylphenyl, 2-methylphenyl, 2,6-difluorophenyl, 2,3,4-trifluorophenyl, 2,3,6-trifluorophenyl, 3-chlorophenyl, 3-bromophenyl, 3-iodophenyl, 3-chloro-5-fluorophenyl, 3-methoxy-5-trifluoromethylphenyl, 3-trifluoromethyl-4-fluorophenyl, 5,6,7,8-tetrahydronaphthyl, 2-benzylphenyl, phenyl, 3-trifluoromethylphenyl, 5-isoquinolinyl, 2-(N-methylamino)phenyl, 1-naphthyl, 5-quinolinyl, 5-(3-methyl)isoquinolinyl and 3-nitrophenyl;

R³ is hydrogen;

R⁴ is methyl;

R⁵ is hydrogen;

25 and pharmaceutically acceptable salts thereof.

In a class of the invention are compound of the formula (I), wherein

R¹ is selected from the group consisting of 3-trifluoromethylphenyl, 3,5-dichlorophenyl and 3-tolyl;

30 R² is selected from the group consisting of 3,5-dichlorophenyl, 2-(aminophenyl)-phenyl, 2,6-difluorophenyl, 2,3,6-trifluorophenyl, 5,6,7,8-tetrahydronaphthyl, 2-benzylphenyl, 3-trifluoromethylphenyl, 5-isoquinolyl and 5-

quinolinyl;

R³ is hydrogen;

R⁴ is methyl;

R⁵ is hydrogen;

5 and pharmaceutically acceptable salts thereof.

Particularly preferred are compounds of the formula (I) wherein R¹ is 3-trifluoromethylphenyl; R² is selected from 3,5-dichlorophenyl or 5-isoquinolinyl; R³ is hydrogen; R⁴ is methyl; R⁵ is hydrogen; and pharmaceutically acceptable
10 salts thereof.

Illustrative of the invention is a pharmaceutical composition comprising a pharmaceutically acceptable carrier and any of the compounds described above. An illustration of the invention is a pharmaceutical composition made
15 by mixing any of the compounds described above and a pharmaceutically acceptable carrier. Illustrating the invention is a process for making a pharmaceutical composition comprising mixing any of the compounds described above and a pharmaceutically acceptable carrier.

20 Exemplifying the invention is a method of treating a condition mediated by the NPY Y5 receptor in a subject in need thereof comprising administering to the subject a therapeutically effective amount of any of the compounds or pharmaceutical compositions described above.

25 An example of the invention is a method for treating a condition selected from eating disorder, obesity, bulimia nervosa, diabetes, binge eating, anorexia nervosa, dyslipidimia, hypertension, memory loss, epileptic seizures, migraine, sleep disturbances, pain, sexual/reproductive disorders, depression, anxiety, cerebral hemorrhage, shock, congestive heart failure, nasal congestion or
30 diarrhea in a subject in need thereof comprising administering to the subject an effective amount of any of the compounds or pharmaceutical compositions described above.


Further illustrating the invention is the use of a compound of formula I in the preparation of a medicament for treating conditions mediated by the NPY Y5 receptor.

5

DETAILED DESCRIPTION OF THE INVENTION

The present invention provides pyrazole carboxamide derivative compounds, useful as ligands of the neuropeptide Y, subtype 5 receptor. More particularly, the present invention is directed to compounds of the general formula (I):

10



(I)

wherein

R¹, R², R³, R⁴ and R⁵ are as previously defined, and pharmaceutically acceptable salts thereof.

15

The compounds of the present invention may also be present in the form of pharmaceutically acceptable salts. For use in medicine, the salts of the compounds of this invention refer to non-toxic "pharmaceutically acceptable salts." Other salts may, however, be useful in the preparation of compounds according to this invention or of their pharmaceutically acceptable salts. Representative organic or inorganic acids include, but are not limited to, hydrochloric, hydrobromic, hydriodic, perchloric, sulfuric, nitric, phosphoric, acetic, propionic, glycolic, lactic, succinic, maleic, fumaric, malic, tartaric, citric, benzoic, mandelic, methanesulfonic, hydroxyethanesulfonic, benzenesulfonic, oxalic, pamoic, 2-naphthalenesulfonic, p-toluenesulfonic, cyclohexanesulfamic, salicylic, saccharinic or trifluoroacetic acid.

20

25

As used herein, unless otherwise noted, the term "halogen" shall include chlorine, fluorine, bromine and iodine.

- 5 As used herein, unless otherwise noted, the terms "alkyl" and "alkoxy" whether used alone or as part of a substituent group, include straight and branched chains having 1-8 carbon atoms, or any number within this range. For example, alkyl radicals include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, t-butyl, n-pentyl, 2-methyl-3-butyl, n-hexyl and the like.
- 10 Alkoxy radicals are oxygen ethers formed from the previously described straight or branched chain alkyl groups. The terms "fluorinated alkyl" and "fluorinated alkoxy" as used herein refer to an alkyl or alkoxy group wherein one or more of the hydrogen atoms are replaced with a fluorine (e. g. trifluoromethyl, trifluoromethoxy).

15

As used herein, unless otherwise noted, "cycloalkyl" shall include saturated C₃-C₈ ring structures, preferably C₅-C₈ ring structures, for example cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and the like.

- 20 As used herein, unless otherwise noted, "heterocycloalkyl" shall include saturated C₅-C₆ ring structures consisting of carbon and one to three heteroatoms (preferably one or two heteroatoms) selected from N, O or S (preferably N or O). Examples of suitable heterocycloalkyl groups include piperidinyl, morpholino, piperazinyl and the like.

25

- As used herein, unless otherwise noted, "aryl" shall include aromatic groups such as: (a) phenyl, naphthyl, fluorenyl, and the like; (b) partially unsaturated C₉-C₁₀ fused ring systems consisting of a phenyl fused to a five or six membered cycloalkyl (e. g. tetrahydronaphthyl, indanyl) or heterocycloalkyl (e. g. methylenedioxyphenyl, ethylenedioxyphenyl, tetrahydroisoquinoliny) group; and (c) stable unsubstituted or substituted fourteen membered benzo-fused tricyclic ring system (e. g. anthraquinoliny).
- 30


As used herein, unless otherwise noted, "heteroaryl" shall denote: (a) a stable unsubstituted or substituted five or six membered monocyclic aromatic ring system; (b) a stable unsubstituted or substituted nine or ten membered benzo-fused heteroaromatic ring system which consists of carbon atoms and from one to six heteroatoms (preferably, one to four heteroatoms) selected from N, O or S; (c) a stable, unsubstituted or substituted nine or ten membered bicyclic fused heteroaromatic ring system which consists of carbon atoms and from one to six heteroatoms (preferably, one to four heteroatoms) selected from N, O or S (preferably N) or (d) a stable, unsubstituted or substituted fourteen membered benzo-fused tricyclic ring system which consists of carbon atoms and from one to six heteroatoms (preferably, one to three heteroatoms) selected from N, O or S (preferably O). The heteroaryl group may be attached at any heteroatom or carbon atom which results in the creation of a stable structure. Examples of suitable heteroaryl groups include, but are not limited to pyridyl, pyrazinyl, pyridazinyl, pirimidyl, thiophenyl, furanyl, imidazolyl, isoxazolyl, indolyl, indazolyl, isoindolyl, pyrazolyl, pyrrolyl, thiazolyl, thiadiazolyl, oxazolyl, triazolyl, tetrazolyl, purinyl, benzimidazolyl, benzofuranyl, benzothienyl, benzisoxazolyl, benzoxazolyl, benzopyrazolyl, indolyl, benzothiazolyl, benzothiadiazolyl, benzotriazolyl, quinolinyl or isoquinolinyl.

As used herein, unless otherwise noted, "aralkyl" shall mean any alkyl group substituted with an aryl group such as benzyl, phenylethyl and the like. Similarly, the term "aralkoxy" indicates an alkoxy group substituted with an aryl group. The term "aminoalkyl" refers to an alkyl group substituted with an amino group (i.e. -alkyl-NH₂). The term "alkylamino" refers to an amino group substituted with an alkyl group (i.e. -NH-alkyl). The term "dialkylamino" refers to an amino group which is disubstituted with alkyl groups wherein the alkyl group can be the same or different (i.e. -N-[alkyl]₂). Suitable alkyl and aryl groups are as defined above.

As used herein, unless otherwise noted, the term "amido" refers to

–C(O)–NH₂. N-alkylamido and N,N-dialkylamido refers to –C(O)–NH-alkyl and –C(O)–N(alkyl)₂, respectively. Similarly, sulfonamido refers to –SO₂–NH₂.

- Under standard nomenclature used throughout this disclosure, the terminal portion of the designated side chain is described first, followed by the adjacent functionality toward the point of attachment. Thus, for example, a “phenylC₁–C₆ alkylamidoC₁–C₆alkyl” substituent refers to a group of the formula



- When a particular group (e.g. alkyl, aryl, heteroaryl) is substituted, that group may have one or more substituents (preferably, one to five, more preferably, one to three, most preferably, one to two substituents) independently selected from the list of substituents.
- Whenever the term “alkyl” or “aryl” or either of their prefix roots appear in a name of a substituent (e. g. aralkyl, dialkylamino) it shall be interpreted as including those limitations given above for “alkyl” and “aryl”. Designated numbers of carbon atoms (e. g. C₁–C₆) shall refer independently to the number of carbon atoms in an alkyl or cycloalkyl moiety or to the alkyl portion of a larger substituent in which alkyl appears as its prefix root.

- Where the compounds according to this invention have at least one chiral center, they may accordingly exist as enantiomers. Where the compounds possess two or more chiral centers, they may additionally exist as diastereomers. It is to be understood that all such isomers and mixtures thereof are encompassed within the scope of the present invention. Furthermore, some of the crystalline forms for the compounds may exist as polymorphs and as such are intended to be included in the present invention. In addition, some of the compounds may form solvates with water (i.e.,

hydrates) or common organic solvents, and such solvates are also intended to be encompassed within the scope of this invention.

It is intended that the definition of any substituent or variable at a particular location in a molecule be independent of its definitions elsewhere in that molecule. It is understood that substituents and substitution patterns on the compounds of this invention can be selected by one of ordinary skill in the art to provide compounds that are chemically stable and that can be readily synthesized by techniques known in the art as well as those methods set forth herein.

The term "subject" as used herein, refers to an animal, preferably mammal, most preferably a human, who has been the object of treatment, observation or experiment.

15

The term "therapeutically effective amount" as used herein, means that amount of active compound or pharmaceutical agent that elicits the biological or medicinal response in a tissue system, animal or human that is being sought by a researcher, veterinarian, medical doctor or other clinician, which includes alleviation of the symptoms of the disease or disorder being treated.

20

As used herein, the term "composition" is intended to encompass a product comprising the specified ingredients in the specified amounts, as well as any product which results, directly or indirectly, from combinations of the specified ingredients in the specified amounts.

25

For use in medicine, the salts of the compounds of this invention refer to non-toxic "pharmaceutically acceptable salts." Other salts may, however, be useful in the preparation of compounds according to this invention or of their pharmaceutically acceptable salts. Suitable pharmaceutically acceptable salts of the compounds include acid addition salts which may, for example, be formed by mixing a solution of the compound with a solution of a

30

pharmaceutically acceptable acid such as hydrochloric acid, sulfuric acid, fumaric acid, maleic acid, succinic acid, acetic acid, benzoic acid, citric acid, tartaric acid, carbonic acid or phosphoric acid. Furthermore, where the compounds of the invention carry an acidic moiety, suitable pharmaceutically acceptable salts thereof may include alkali metal salts, e.g., sodium or potassium salts; alkaline earth metal salts, e.g., calcium or magnesium salts; and salts formed with suitable organic ligands, e.g., quaternary ammonium salts. Thus, representative pharmaceutically acceptable salts include the following:

10 acetate, benzenesulfonate, benzoate, bicarbonate, bisulfate, bitartrate, borate, bromide, calcium edetate, camsylate, carbonate, chloride, clavulanate, citrate, dihydrochloride, edetate, edisylate, estolate, esylate, fumarate, gluceptate, gluconate, glutamate, glycolylarsanilate, hexylresorcinate, hydrabamine, hydrobromide, hydrochloride, hydroxynaphthoate, iodide,

15 isothionate, lactate, lactobionate, laurate, malate, maleate, mandelate, mesylate, methylbromide, methylnitrate, methylsulfate, mucate, napsylate, nitrate, N-methylglucamine ammonium salt, oleate, pamoate (embonate), palmitate, pantothenate, phosphate/diphosphate, polygalacturonate, salicylate, stearate, sulfate, subacetate, succinate, tannate, tartrate, teoclate, tosylate,

20 triethiodide and valerate.

The present invention includes within its scope prodrugs of the compounds of this invention. In general, such prodrugs will be functional derivatives of the compounds which are readily convertible in vivo into the required compound. Thus, in the methods of treatment of the present invention, the term "administering" shall encompass the treatment of the various disorders described with the compound specifically disclosed or with a compound which may not be specifically disclosed, but which converts to the specified compound in vivo after administration to the patient. Conventional procedures for the selection and preparation of suitable prodrug derivatives are described, for example, in "Design of Prodrugs", ed. H. Bundgaard, Elsevier, 1985.

25


30

The compounds of formula (I) that comprise this invention are generally referred to as pyrazole carboxamide derivatives, and are synthesized via the route outlined in Scheme 1; consisting of several sequential chemical

5 operations that can be generalized as described below:

- Formation of the pyrazole nucleus (ring closure)
- Hydrolysis of the carboxylic ester to the carboxylic acid
- Coupling of the acid to an appropriate amine


10 In general, the synthesis of compounds of formula (I) consists of the steps of reacting a diketoeester of formula (II) with an aryl or heteroaryl hydrazine to produce the 1,3,5-trisubstituted pyrazole of formula (III), further reacting the compound of formula (III) with a base to yield the corresponding carboxylic acid of formula (IV) and further reacting the carboxylic acid of
15 formula (IV) with an aryl, heteroaryl or alkyl amine to afford the pyrazole carboxamide derivative of formula (I).



SCHEME 1

Specifically, a diketoester of the formula (II) is reacted with an aryl or heteroaryl hydrazine in the presence of an acid such as acetic acid, hydrochloric acid and the like, wherein the reaction solution is heated from ambient temperature to reflux, to afford the corresponding substituted pyrazoles of formula (III). The pyrazole of formula (III) is hydrolyzed in the presence of a base, such as sodium hydroxide, potassium carbonate and the like, in an aqueous alcoholic solvent, such as an aqueous methanolic solution, aqueous ethanolic solution and the like, wherein the reaction solution is heated from ambient temperature to reflux, to yield the pyrazole carboxylic acid of formula (IV). The pyrazole carboxylic acid of formula (IV) is coupled to an aryl, heteroaryl or alkyl amine (which may be primary or secondary) in the presence of a sterically hindered, non-nucleophilic amine, such as diisopropyl ethyl amine, triethyl amine and the like, and a coupling agent, such as O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU), to provide the pyrazole carboxamides of formula (I) (Scheme 1).

The substituents R^3 and R^4 are varied by methods known to those skilled in the art, such as acylation of an appropriately substituted ketone of formula (V) with diethyl oxalate, in the presence of a base such as sodium hydride, sodium t-butoxide, lithium diisopropyl amide, and the like, to form the diketoester of formula (II), as shown in Scheme II.




SCHEME II

Alternately, the R^3 and R^4 substituents may be introduced via a process

comprising the steps of acylating an appropriately substituted methyl ketone of formula (VI), with diethyl oxalate, in the presence of a base such as sodium hydride, sodium t-butoxide, lithium diisopropyl amide, and the like, to form the diketoester of formula (VII),

5



SCHEME III

reacting the compound of formula (VII) (to introduce R³) to form the compound of formula (II), via synthesis routes known to those skilled in the art, such as alkylation, electrophilic halogenation, displacement reactions of intermediate halo species, electrophilic amination and the like.

Those compounds of the present invention which contain a basic moiety can be converted to the corresponding acid addition salts by techniques known to those skilled in the art. Suitable acids which can be employed for this purpose include hydrochloric, hydrobromic, hydroiodic, perchloric, sulfuric, nitric, phosphoric, acetic, propionic, glycolic, lactic, pyruvic, oxalic, malonic, succinic, maleic, fumaric, malic, tartaric, citric, benzoic, cinnamic, mandelic, methanesulfonic, p-toluenesulfonic, cyclohexanesulfamic, salicylic, 2-phenoxybenzoic, 2-acetoxybenzoic, or saccharin, and the like. In general, the acid addition salts can be prepared by reacting the free base of compounds of formula (I) with the acid and isolating the salt.

It is generally preferred that the respective product of each process step be separated from other components of the reaction mixture and then subjected to purification before its use as a starting material in a subsequent step. Separation techniques typically include evaporation, extraction, precipitation and filtration. Purification techniques typically include column

chromatography (Still, W. C. et. al., J. Org. Chem. 1978, 43, 2921), thin-layer chromatography, crystallization and distillation. In those cases wherein the product is isolated as the acid addition salt, the free base is obtained by techniques known to those skilled in the art. The structures of the final products, intermediates and starting materials are confirmed by spectroscopic, spectrometric and analytical methods including nuclear magnetic resonance (NMR), mass spectrometry (MS) and liquid chromatography (HPLC).

During any of the processes for preparation of the compounds of the present invention, it may be necessary and/or desirable to protect sensitive or reactive groups on any of the molecules concerned. This may be achieved by means of conventional protecting groups, such as those described in Protective Groups in Organic Chemistry, ed. J.F.W. McOmie, Plenum Press, 1973; and T.W. Greene & P.G.M. Wuts, Protective Groups in Organic Synthesis, John Wiley & Sons, 1991. The protecting groups may be removed at a convenient subsequent stage using methods known from the art.

As modulators of the NPY5 receptor, the compounds of Formula (I) are useful for treating feeding disorders such as obesity, binge eating, anorexia nervosa and bulimia nervosa, and abnormal conditions such as epilepsy, depression, anxiety and sexual / reproductive disorders in which modulation of the NPY5 receptor may be useful. The compounds compete with the endogenous ligands NPY and PYY and possibly non-endogenous ligands, and bind to the NPY5 receptor. In addition, the compounds demonstrate antagonist activity by antagonizing the action of NPY upon binding to the Y5 receptor.

The compounds described herein are ligands of the NPY5 receptor, but are not necessarily limited solely in their pharmacological or biological action due to binding to this or any neuropeptide, neurotransmitter or G-protein coupled receptor. For example, the described compounds may also undergo binding to dopamine or serotonin receptors. The compounds described herein

are potentially useful in the regulation of metabolic and endocrine functions, particularly those associated with feeding, and as such, may be useful for the treatment of obesity. In addition, the compounds described herein are potentially useful for modulating other endocrine functions, particularly those controlled by the pituitary and hypothalamic glands, and therefore may be useful for the treatment of ovulation/infertility due to insufficient release of luteinizing hormone (LH).

The present invention further comprises pharmaceutical compositions containing one or more compounds of formula (I). The pharmaceutical composition of this invention alternately comprise a pharmaceutically acceptable carrier and one or more of the compounds of formula (I) described above. Pharmaceutical compositions containing one or more of the compounds of the invention described herein as the active ingredient can be prepared by intimately mixing the compound or compounds with a pharmaceutical carrier according to conventional pharmaceutical compounding techniques. The carrier may take a wide variety of forms depending upon the desired route of administration (e.g., oral, parenteral). Thus for liquid oral preparations such as suspensions, elixirs and solutions, suitable carriers and additives include water, glycols, oils, alcohols, flavoring agents, preservatives, stabilizers, coloring agents and the like; for solid oral preparations, such as powders, capsules and tablets, suitable carriers and additives include starches, sugars, diluents, granulating agents, lubricants, binders, disintegrating agents and the like. Solid oral preparations may also be coated with substances such as sugars or be enteric-coated so as to modulate major site of absorption. For parenteral administration, the carrier will usually consist of sterile water and other ingredients may be added to increase solubility or preservation. Injectable suspensions or solutions may also be prepared utilizing aqueous carriers along with appropriate additives.

30

To prepare the pharmaceutical compositions of this invention, one or more compounds of formula (I) or salt thereof of the invention as the active

ingredient, is intimately admixed with a pharmaceutical carrier according to conventional pharmaceutical compounding techniques, which carrier may take a wide variety of forms depending of the form of preparation desired for administration, e.g., oral or parenteral such as intramuscular. In preparing the compositions in oral dosage form, any of the usual pharmaceutical media may be employed. Thus, for liquid oral preparations, such as for example, suspensions, elixirs and solutions, suitable carriers and additives include water, glycols, oils, alcohols, flavoring agents, preservatives, coloring agents and the like; for solid oral preparations such as, for example, powders, capsules, caplets, gelcaps and tablets, suitable carriers and additives include starches, sugars, diluents, granulating agents, lubricants, binders, disintegrating agents and the like. Because of their ease in administration, tablets and capsules represent the most advantageous oral dosage unit form, in which case solid pharmaceutical carriers are obviously employed. If desired, tablets may be sugar coated or enteric coated by standard techniques. For parenterals, the carrier will usually comprise sterile water, through other ingredients, for example, for purposes such as aiding solubility or for preservation, may be included. Injectable suspensions may also be prepared, in which case appropriate liquid carriers, suspending agents and the like may be employed. The pharmaceutical compositions herein will contain, per dosage unit, e.g., tablet, capsule, powder, injection, teaspoonful and the like, an amount of the active ingredient necessary to deliver an effective dose as described above. The pharmaceutical compositions herein will contain, per unit dosage unit, e.g., tablet, capsule, powder, injection, suppository, teaspoonful and the like, of from about 0.03 mg to 500 mg/kg (preferred 0.1-150 mg/kg) and may be given at a dosage of from about 0.1-300 mg/kg/day (preferred 1-150 mg/kg/day). The dosages, however, may be varied depending upon the requirement of the patients, the severity of the condition being treated and the compound being employed. The use of either daily administration or post-periodic dosing may be employed.

Preferably these compositions are in unit dosage forms from such as

tablets, pills, capsules, powders, granules, sterile parenteral solutions or suspensions, metered aerosol or liquid sprays, drops, ampoules, autoinjector devices or suppositories; for oral parenteral, intranasal, sublingual or rectal administration, or for administration by inhalation or insufflation. Alternatively, the composition may be presented in a form suitable for once-weekly or once-monthly administration; for example, an insoluble salt of the active compound, such as the decanoate salt, may be adapted to provide a depot preparation for intramuscular injection. For preparing solid compositions such as tablets, the principal active ingredient is mixed with a pharmaceutical carrier, e.g. conventional tableting ingredients such as corn starch, lactose, sucrose, sorbitol, talc, stearic acid, magnesium stearate, dicalcium phosphate or gums, and other pharmaceutical diluents, e.g. water, to form a solid preformulation composition containing a homogeneous mixture of a compound of the present invention, or a pharmaceutically acceptable salt thereof. When referring to these preformulation compositions as homogeneous, it is meant that the active ingredient is dispersed evenly throughout the composition so that the composition may be readily subdivided into equally effective dosage forms such as tablets, pills and capsules. This solid preformulation composition is then subdivided into unit dosage forms of the type described above containing from 0.1 to about 500 mg of the active ingredient of the present invention. The tablets or pills of the novel composition can be coated or otherwise compounded to provide a dosage form affording the advantage of prolonged action. For example, the tablet or pill can comprise an inner dosage and an outer dosage component, the latter being in the form of an envelope over the former. The two components can be separated by an enteric layer which serves to resist disintegration in the stomach and permits the inner component to pass intact into the duodenum or to be delayed in release. A variety of material can be used for such enteric layers or coatings, such materials including a number of polymeric acids with such materials as shellac, cetyl alcohol and cellulose acetate.

The liquid forms in which the novel compositions of the present

invention may be incorporated for administration orally or by injection include, aqueous solutions, suitably flavoured syrups, aqueous or oil suspensions, and flavoured emulsions with edible oils such as cottonseed oil, sesame oil, coconut oil or peanut oil, as well as elixirs and similar pharmaceutical vehicles.

5 Suitable dispersing or suspending agents for aqueous suspensions, include synthetic and natural gums such as tragacanth, acacia, alginate, dextran, sodium carboxymethylcellulose, methylcellulose, polyvinyl-pyrrolidone or gelatin.

10 The present invention further provides a method of treatment of central nervous system disorders and affective conditions such as eating disorder, obesity, bulimia nervosa, diabetes, binge eating, anorexia nervosa, dyslipidimia, hypertension, memory loss, epileptic seizures, migraine, sleep disturbances, pain, sexual/reproductive disorders, depression, anxiety,
15 cerebral hemorrhage, shock, congestive heart failure, nasal congestion or diarrhea.

 The utility of the compounds to treat disorders of the central nervous system as described above, can be determined according to the procedures
20 described herein. The present invention therefore provides a method of treating central nervous system disorders in a subject in need thereof which comprises administering any of the compounds as defined herein in a quantity effective to treat central nervous system disorders. The compound may be administered to a patient by any conventional route of administration,
25 including, but not limited to, intravenous, oral, subcutaneous, intramuscular, intradermal and parenteral.

 The method of treating central nervous system disorders described in the present invention may also be carried out using a pharmaceutical composition
30 comprising any of the compounds as defined herein and a pharmaceutically acceptable carrier. The pharmaceutical composition may contain between about 0.01 mg and 500 mg, preferably about 5 to 150 mg, of the compound, and may

be constituted into any form suitable for the mode of administration selected. Carriers include necessary and inert pharmaceutical excipients, including, but not limited to, binders, suspending agents, lubricants, flavorants, sweeteners, preservatives, dyes, and coatings. Compositions suitable for oral administration include solid forms, such as pills, tablets, caplets, capsules (each including immediate release, timed release and sustained release formulations), granules, and powders, and liquid forms, such as solutions, syrups, elixers, emulsions, and suspensions. Forms useful for parenteral administration include sterile solutions, emulsions and suspensions.

10

Advantageously, compounds of the present invention may be administered in a single daily dose, or the total daily dosage may be administered in divided doses of two, three or four times daily. Furthermore, compounds for the present invention can be administered in intranasal form via topical use of suitable intranasal vehicles, or via transdermal skin patches well known to those of ordinary skill in that art. To be administered in the form of a transdermal delivery system, the dosage administration will, of course, be continuous rather than intermittent throughout the dosage regimen.

15

20

For instance, for oral administration in the form of a tablet or capsule, the active drug component can be combined with an oral, non-toxic pharmaceutically acceptable inert carrier such as ethanol, glycerol, water and the like. Moreover, when desired or necessary, suitable binders; lubricants, disintegrating agents and coloring agents can also be incorporated into the mixture. Suitable binders include, without limitation, starch, gelatin, natural sugars such as glucose or beta-lactose, corn sweeteners, natural and synthetic gums such as acacia, tragacanth or sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride and the like. Disintegrators include, without limitation, starch, methyl cellulose, agar, bentonite, xanthan gum and the like.

25

30

The liquid forms in suitably flavored suspending or dispersing agents

such as the synthetic and natural gums, for example, tragacanth, acacia, methyl-cellulose and the like. For parenteral administration, sterile suspensions and solutions are desired. Isotonic preparations which generally contain suitable preservatives are employed when intravenous administration is desired.

5

The compound of the present invention can also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles, and multilamellar vesicles. Liposomes can be formed from a variety of phospholipids, such as cholesterol, stearylamine or

10 phophatidylcholines.

Compounds of the present invention may also be delivered by the use of monoclonal antibodies as individual carriers to which the compound molecules are coupled. The compounds of the present invention may also be coupled with

15 soluble polymers as targetable drug carriers. Such polymers can include polyvinylpyrrolidone, pyran copolymer, polyhydroxypropylmethacrylamidephenol, polyhydroxy-ethylaspartamidephenol, or polyethyl eneoxydepolylysine substituted with palmitoyl residue. Furthermore, the compounds of the present invention may be coupled to a class of biodegradable polymers useful in

20 achieving controlled release of a drug, for example, polylactic acid, polyepsilon caprolactone, polyhydroxy butyeric acid, polyorthoesters, polyacetals, polydihydropyrans, polycyanoacrylates and cross-linked or amphipathic block copolymers of hydrogels.

25 Compounds of this invention may be administered in any of the foregoing compositions and according to dosage regimens established in the art whenever treatment of disorders of the central nervous system is required.

The daily dosage of the products may be varied over a wide range from

30 0.01 to 1,000 mg per adult human per day. For oral administration, the compositions are preferably provided in the form of tablets containing, 0.01, 0.05, 0.1, 0.5, 1.0, 2.5, 5.0, 10.0, 15.0, 25.0, 50.0, 100, 150, 200, 250 and 500

milligrams of the active ingredient for the symptomatic adjustment of the dosage to the patient to be treated. An effective amount of the drug is ordinarily supplied at a dosage level of from about 0.01 mg/kg to about 500 mg/kg of body weight per day. Preferably, the range is from about 0.03 to about 300 mg/kg of body weight per day. The compounds may be administered on a regimen of 1 to 4 times per day.

Optimal dosages to be administered may be readily determined by those skilled in the art, and will vary with the particular compound used, the mode of administration, the strength of the preparation, the mode of administration, and the advancement of the disease condition. In addition, factors associated with the particular patient being treated, including patient age, weight, diet and time of administration, will result in the need to adjust dosages.

The following example(s) describe the invention in greater detail and are intended to illustrate the invention, but not to limit it. All compounds were identified by a variety of methods including nuclear magnetic resonance spectroscopy, mass spectrometry and in some cases, infrared spectroscopy and elemental analysis. Nuclear magnetic resonance (300 MHz NMR) data is reported in parts per million downfield from tetramethylsilane. Mass spectra data is reported in mass/charge (m/z) units. Unless otherwise noted, the materials used in the example were obtained from readily available commercial sources or synthesized by standard methods known to those skilled in the art.

Abbreviations used in the instant specification, particularly the Schemes and Examples, are as follows:

BSA	=	Bovine Serum Albumin
Cmpd	=	Compound
DIPEA	=	Diisopropyl Ethyl Amine
EDTA	=	Ethylene Diamine Tetraacetic Acid
EtOAc	=	Ethyl acetate

HATU	=	O-(7-Azabenzotriazol-1yl)-N,N,N",N"-Tetramethyl Uronium Hexafluorophosphate
HEPES	=	4-(2-hydroxyethyl)-1-Piperazine Ethane Sulfonic Acid
HPLC	=	High Performance Liquid Chromatography
% Inh	=	Percent Inhibition
MeOH	=	Methanol
PBS	=	Phosphate Buffered Saline
PEG	=	Poly(Ethylene) Glycol
RT or rt	=	Room temperature

EXAMPLE 1

1-[(3-Trifluoromethyl)phenyl]-3-[N-5-(isoquinolinyl)carboxamide)]-5-methylpyrazole (110).

- 5 A. To a solution of ethyl 2,4-dioxovalerate (2.5 g, 16.0 mmol) in a 2:1 mixture of acetic acid and 2-methoxyethanol (48 mL) at room temperature in a 250 mL round-bottom flask fitted with a reflux condenser under nitrogen was added 3-(trifluoromethyl)phenylhydrazine (3.3g, 19.0 mmol). The resultant solution was heated to reflux for 18 h. After cooling to room temperature,
- 10 the volatiles were removed in vacuo and the residue was partitioned between EtOAc (200 mL) and 1N aq. HCl solution (200 mL). The layers were separated and the organic layer was washed with H₂O (100 mL), dried over Na₂SO₄, and concentrated. The residue was purified by column chromatography on silica gel eluting with 5-10% EtOAc/hexanes provide
- 15 the pyrazole ester III [R₁ = 3-(trifluoromethyl)phenyl, R₃ = H, R₄ = CH₃]. ¹H NMR (CDCl₃, 300 MHz) δ 1.42 (t, 3H, J = 8.6 Hz), 2.40 (s, 3H), 4.42 (q, 2H, J = 8.6 Hz), 6.78 (s, 1H), 7.65 (m, 3H), 7.79 (s 1H).
- B. To a solution of pyrazole carboxylic ester III [R₁ = 3-(trifluoromethyl)phenyl; 2.19 g, 7.3 mmol] in a 3:1 MeOH/H₂O solution (150 mL) at room
- 20 temperature in a 500 mL round-bottom flask fitted with a reflux condenser under nitrogen was added NaOH (440 mg, 11.0 mmol). The solution was heated to reflux for 18 h. The solution was allowed to cool to room

temperature and was acidified to a pH of 2 with concentrated HCl. The solution was concentrated and the residue partitioned between CH₂Cl₂ (200 mL) and H₂O (200 mL). The layers were separated and the aqueous layer was extracted with 2 X 200 mL CH₂Cl₂. The combined organic layers were dried over Na₂SO₄ and concentrated to provide the acid IV [R₁ = 3-(trifluoromethyl)phenyl, R₃ = H, R₄ = CH₃]. ¹H NMR (CDCl₃, 300 MHz) δ 2.41 (s, 3H), 6.84 (s, 1H), 7.70 (m, 3H), 7.78 (s, 1H).

5 C. To a solution of pyrazole carboxylic acid IV [R₁ = 3-(trifluoromethyl)phenyl; 100 mg, 0.37 mmol] in CH₂Cl₂ (5 mL) at room temperature was added DIPEA (0.13 mL) and HATU (142 mg, 0.37 mmol). The solution was allowed to stir at room temperature for 10 min. and then to it was added 5-aminoisoquinoline (59 mg, 0.41 mmol). The reaction mixture was allowed to stir at room temperature overnight, and then diluted with H₂O (10 mL) and extracted 3 X 10 mL of CH₂Cl₂. The organic layers were combined and washed with 2 X 10 mL 1N aq. HCl, 2 X 10 mL H₂O, dried over Na₂SO₄, and concentrated. The residue was purified by column chromatography eluting with 10% EtOAc/hexanes to provide the pyrazole carboxamide I [R₁ = 3-(trifluoromethyl)phenyl, R₂ = 5-quinolinyl, R₃ = H, R₄ = CH₃, Compound 110]. ¹H NMR (CDCl₃, 300 MHz) δ 2.80 (s, 3H), 6.90 (s, 1H), 7.60 - 7.87 (m, 6H), 8.46 (d, 1H, J = 8.5 Hz), 8.55 (d, 1H, J = 6.8 Hz), 9.25 (s, 1H), 9.31 (s, 1H).

In a similar manner, all of the compounds of the present invention were been prepared by varying the necessary hydrazines and anilines.

EXAMPLE 2

As a specific embodiment of an oral composition, 100 mg of the compound 110 Of Example 1 is formulated with sufficient finely divided lactose to provide a total amount of 580 to 590 mg to fill a size O hard gel capsule.

EXAMPLE 3

IN VITRO ASSAYS

NPY5 HTS Centrifugation Assay

The compounds described in this invention were evaluated for binding to the human neuropeptide Y5 receptor.

5

Stable Transfection

The human NPY5 receptor cDNA (Genbank Accession number U66275) was inserted into the vector pCIneo (Invitrogen) and transfected into human embryonic kidney cells (HEK-293) via Calcium phosphate method (Cullen 1987). Stably transfected cells were selected with G-418 (600 ug/mL). Stably transfected cells served as the source for the membranes for the NPY5 receptor binding assay.

Membrane Preparation

15 PY5-transfected HEK293 cells were grown to confluence in 150 cm² culture dishes. Cells were washed once with phosphate-buffered saline (Gibco Cat# 14040-133). Cells were then incubated in phosphate-buffered saline without Calcium and without Magnesium, supplemented with 2 mM EDTA. Cells were incubated for 10 minutes at room temperature and the cells were collected by repetitive pipeting. Cells were formed into pellets and then frozen at -80 until needed. Frozen pellets were homogenized with a polytron at full speed for 12 seconds in a homogenization buffer (20 mM Tris HCl, 5 mM EDTA, pH 7.4). Homogenates were centrifuged for 5 minutes at 4C at 200g. Supernatants were transferred to corex tubes and centrifuged for 25 minutes at 28,000g. Pellets were re-suspended in Binding (20mM HEPES, 10 mM NaCl, 0.22 mM KH₂PO₄, 1.3mM CaCl₂, 0.8 mM MgSO₄, pH 7.4). Membranes were kept on ice until use.


30 A competition binding assay, known to those skilled in the art, was used in which the compounds of formula (I) compete with ¹²⁵I-PYY for binding to cell membranes. In simple terms, the less ¹²⁵I-PYY bound to the membranes implies that a compound is a good inhibitor (competitor). Bound ¹²⁵I-PYY was

determined by centrifugation of membranes, aspirating supernatant, washing away residual ^{125}I -PYY and subsequently counting the bound sample in a g-counter.

5 Procedure for Radioligand binding assay



Compounds to be tested were prepared as 10x stocks in binding buffer and added first to assay tubes (RIA vials, Sarstedt). Twenty (20) μL of each 10x compound stock was pipeted into vials and 80 μL of ^{125}I -PYY (NEN catalog number NEX240), which had been diluted to a concentration of 200 pM in 0.25
10 % BSA in binding buffer, was added to the compound tubes (final concentration of ^{125}I -PYY is 80 pM). To each tube was added 100 μL of membranes and the mixture was agitated by pipeting 2 times. Samples were incubated for 1 hr at room temperature. Aluminum cast plates (Sarstedt) containing the vials were then centrifuged 10 minutes at 3200 rpm in a Sorvall
15 RT6000. Supernatant was then aspirated. To each vial 400 μL PBS was added and this was then aspirated again. Vials were then put in carrier polypropylene 12x75 tube and counted in gamma counter (Packard). Non-specific binding was determined in the presence of 300 nM NPY. Percent inhibition of ^{125}I -PYY binding was calculated by subtracting non-specific binding
20 from the test samples (compound (I)), taking these counts and dividing by total binding, and multiplying by 100. Inhibitory concentration values (IC_{50}) of compounds that showed appreciable inhibition of ^{125}I -PYY binding were calculated by obtaining percent inhibition of ^{125}I -PYY binding values at different concentrations of the test compound, and using a graphing program such as
25 GraphPad Prism (San Diego, CA) the concentration of test compound that inhibits fifty-percent of ^{125}I -PYY binding (Table 4) was calculated. These operations are known to those skilled in the art.



Mass Spectral Data and
Binding Affinities of Compounds (I) for the Human NPY Y5 Receptor
(expressed as % Inhibition of 125 I-PYY Binding or IC_{50})








5




R^3 = hydrogen, R^4 = methyl, R^5 = hydrogen

Cmpd #	R^1	R^2	%Inh NPY5r 3 μ M	IC_{50} HNPY5r	Mass (calcd)	Mass (obs)
1	4-tolyl	4-fluorophenyl	36	8 μ M	309.34	310.3
2	4-tolyl	3,5-dichlorophenyl	67	3 μ M	360.24	360
3	4-tolyl	4-methoxyphenyl	7	20 μ M	321.38	322
4	4-tolyl	4-chlorophenyl	29		325.80	326
5	3-trifluoromethyl phenyl	4-fluorophenyl	86	828 nM	363.31	364
6	3-trifluoromethyl phenyl	3,5-dichlorophenyl	102	124 nM	414.21	414.05
7	3-trifluoromethyl phenyl	4-methoxyphenyl	34		375.35	376
8	3-trifluoromethyl phenyl	4-(N-morpholino)- phenyl	8		430.43	431
9	3-trifluoromethyl phenyl	4-chlorophenyl	48		379.77	380
10	3-trifluoromethyl phenyl	2-(1-pyrrolyl) phenyl	75	763 nM	410.40	411
11	3-trifluoromethyl phenyl		20		442.48	443
12	3-trifluoromethyl phenyl		30		367.37	368











13	3-trifluoromethyl phenyl		22		408.47	409
14	3-trifluoromethyl phenyl	$-(CH_2)_3N(CH_3)_2$	16		354.37	355
15	3-trifluoromethyl phenyl	-2-(aminophenyl) phenyl	91	267 nM	436.44	437.14
16	3-trifluoromethyl phenyl		0		461.45	462.16
17	3-trifluoromethyl phenyl	2-chlorophenyl	2		379.77	380.36
18	3-trifluoromethyl phenyl	2-bromophenyl	3		424.22	425.35
19	3-trifluoromethyl phenyl	2-iodophenyl	10		471.22	472.46
20	3-trifluoromethyl phenyl	4-iodophenyl	0		471.22	472.27
21	3-trifluoromethyl phenyl	3,4-dichlorophenyl	47	1.5 μ M	414.21	415.28
22	3-trifluoromethyl phenyl	3,5- dimethoxyphenyl	48		405.37	406.42
23	3-trifluoromethyl phenyl	3-trifluoromethoxy phenyl	83	431 nM	429.32	430
24	3-trifluoromethyl phenyl	4-trifluoromethoxy phenyl	5		429.32	430
25	3-trifluoromethyl phenyl	2-methylthio phenyl	63	881 nM	391.41	392
26	3-trifluoromethyl phenyl	2-methoxyphenyl	6		375.35	376
27	3-trifluoromethyl phenyl	3-tolyl	78	622 nM	359.35	360
28	3-trifluoromethyl phenyl	2-tolyl	87	573 nM	359.35	360





29	3-trifluoromethyl phenyl	3,5-ditrifluoro methylphenyl	79		481.32	482
30	3-trifluoromethyl phenyl	2,3-difluorophenyl	83		381.30	382
31	3-trifluoromethyl phenyl	2,5-difluorophenyl	25		381.30	382
32	3-trifluoromethyl phenyl	2,6-difluorophenyl	95	233 nM	381.30	382
33	3-trifluoromethyl phenyl	3,4-difluorophenyl	82		381.30	382
34	3-trifluoromethyl phenyl	2,3,4- trifluorophenyl	87	885 nM	399.29	400
35	3-trifluoromethyl phenyl	2,4,5- trifluorophenyl	0		399.29	400
36	3-trifluoromethyl phenyl	2,4,6- trifluorophenyl	12		399.29	400
37	3-trifluoromethyl phenyl	2,3,6- trifluorophenyl	96	229 nM	399.29	400
38	3-trifluoromethyl phenyl	2-fluorophenyl	72		363.31	364.09
39	3-trifluoromethyl phenyl	3-chlorophenyl	93	520 nM	379.77	380.07
40	3-trifluoromethyl phenyl	3-bromophenyl	94	359 nM	424.22	426.04
41	3-trifluoromethyl phenyl	3-iodophenyl	87	670 nM	471.21	472.01
42	3-trifluoromethyl phenyl	2,5- dibromophenyl	5		503.11	504
43	3-trifluoromethyl phenyl	2,4- dibromophenyl	13		503.11	504
44	3-trifluoromethyl phenyl	3-chloro-5- fluorophenyl	95	516 nM	397.76	398
45	3-trifluoromethyl phenyl	2,5- dimethoxyphenyl	37		405.37	406.37
46	3-trifluoromethyl phenyl	3-methoxy-5- trifluoromethyl phenyl	93	346 nM	443.34	444.40




47	3-trifluoromethyl phenyl	3-trifluoromethyl- 4-fluorophenyl	90	593 nM	431.30	431.17
48	3-trifluoromethyl phenyl	3-trifluoromethyl phenyl	78		413.32	413.25
49	3-trifluoromethyl phenyl	2,6-dichlorophenyl	4		414.21	415.22
50	3-trifluoromethyl phenyl		30		389.33	390.40
51	3-trifluoromethyl phenyl		40		403.36	404.41
52	3-trifluoromethyl phenyl	2,4,6-trifluoro methylphenyl	74		387.40	388.36
53	3-trifluoromethyl phenyl		19		428.45	429.20
54	3-trifluoromethyl phenyl	2-biphenyl	94		421.42	422.14
55	3-trifluoromethyl phenyl		96	197 nM	399.41	400.19
56	3-trifluoromethyl phenyl	2-benzylphenyl	93	292 nM	435.45	436.18
57	3-trifluoromethyl phenyl		59		385.39	386.15
58	3-trifluoromethyl phenyl	2-benzoylphenyl	57		449.43	450.15
59	3-trifluoromethyl phenyl	3- benzyloxyphenyl	35		451.45	452.17
60	3-trifluoromethyl phenyl	2,4,5- trichlorophenyl	0		448.66	449









61	4-trifluoromethyl phenyl	4-fluorophenyl	28		363.31	364.1
62	4-trifluoromethyl phenyl	4-chlorophenyl	13		379.77	380.1
63	4-trifluoromethyl phenyl	2-(1-pyrrolyl) phenyl	16		410.40	411.1
64	4-trifluoromethyl phenyl	4-methoxyphenyl	8		375.35	376.1
65	4-trifluoromethyl phenyl	3,5-dichlorophenyl	49		414.22	414
66	4-trifluoromethyl phenyl	4-tolyl	60		514.53	515.3
67	benzyl	4-fluoromethyl phenyl	14		309.34	310.1
68	benzyl	3,5-dichlorophenyl	57		360.24	360.0
69	benzyl	2-(1-pyrrolyl) phenyl	17		356.43	357.2
70	benzyl	4-chlorophenyl	14		325.80	326.1
71	3-trifluoromethyl phenyl	2-pyridyl	45		346.31	347.11
72	3-trifluoromethyl phenyl	3-pyridyl	83		346.31	347.11
73	3-trifluoromethyl phenyl		16		387.32	389.10
74	3-trifluoromethyl phenyl		30		347.30	348.10
75	3-trifluoromethyl phenyl		81		514.53	515.13
76	2-trifluoromethyl phenyl	4-fluorophenyl	13		363.31	364.1
77	2-trifluoromethyl phenyl	4-chloromethyl phenyl	11		379.77	380.1
78	2-trifluoromethyl phenyl	2-(1-pyrrolyl) phenyl	34		410.40	411.1

79	2-trifluoromethyl phenyl	3,5-dichlorophenyl	22		414.21	414.1
80	2-trifluoromethyl phenyl	4-methoxyphenyl	4		375.35	376.2
81	phenyl	3,5-dichlorophenyl	90	759 nM	346.21	346.0
82	phenyl	4-chlorophenyl	28		311.77	312.0
83	phenyl	4-fluorophenyl	39		295.31	296.1
84	phenyl	4-methoxyphenyl	12		307.35	308.1
85	phenyl	2-(1-pyrrolyl) phenyl	21		342.40	343.1
86	phenyl	Phenyl	39		277.32	278.1
87	3-trifluoromethyl phenyl	Phenyl	84	1 μ M	345.32	346.1
88	3,5- dichlorophenyl	4-fluorophenyl	84	2 μ M	364.20	364.0
89	3,5- dichlorophenyl	4-chlorophenyl		6 μ M	380.66	380.0
90	3,5- dichlorophenyl	3,5-dichlorophenyl		1 μ M	415.11	415.9
91	3,5- dichlorophenyl	Phenyl		3 μ M	346.21	346.0
92	3,5- dichlorophenyl	2-(1-pyrrolyl) phenyl		16 μ M	411.29	411.1
93	3-trifluoromethyl phenyl	2,3-dichlorophenyl		1 μ M	414.21	414.1
94	3,5- dichlorophenyl	2,3-dichlorophenyl	0		415.10	416
95	3,5- dichlorophenyl	3,5-trifluoromethyl phenyl		5 μ M	482.21	481.8
96	3,5- dichlorophenyl	3-trifluoromethyl phenyl		261 nM	414.21	414.1
97	3-tolyl	4-fluorophenyl	68		309.34	310.1
98	3-tolyl	4-chlorophenyl	57		325.80	326.1
99	3-tolyl	3,5-dichlorophenyl	93	218 nM	360.24	360.1
100	3-tolyl	Phenyl	54		291.35	292.1
101	3-tolyl	2-(1-pyrrolyl) phenyl	56		356.43	357.2

102	3-trifluoromethyl phenyl		0		420.39	421.0
103	3-trifluoromethyl phenyl			5 μ M	460.46	461.1
104	3-trifluoromethyl phenyl			14 μ M	399.37	400.2
105	3-trifluoromethyl phenyl		0		492.52	493.1
106	3-trifluoromethyl phenyl		0		396.37	397.2
107	3-trifluoromethyl phenyl			4 μ M	396.37	397.2
108	3-trifluoromethyl phenyl			13 μ M	396.37	397.2
109	3-trifluoromethyl phenyl			6 μ M	433.43	434.1
110	3-trifluoromethyl phenyl			80 nM	396.37	397.1
111	3-trifluoromethyl phenyl	3-biphenyl	33		421.42	422.15
112	3-trifluoromethyl phenyl		22		470.53	471.4
113	3-trifluoromethyl phenyl	2-(N-methyl amino)phenyl	95	343 nM	374.36	375.2
114	2-pyridyl	4-fluorophenyl	0		296.30	297.11
115	2-pyridyl	3,5-dichlorophenyl	0		347.20	347.03

116	2-pyridyl	2-(1-pyrrolyl) phenyl	5		343.39	344.14
117	2-pyridyl	2-(N-phenyl amino) phenyl	18		369.42	370.16
118	3-trifluoromethyl phenyl	4-fluorobenzyl	54		377.34	378.12
119	3-trifluoromethyl phenyl	3,5-dichlorobenzyl	38		428.24	428.07 430.07
120	3-trifluoromethyl phenyl	3,5-trifluoromethyl benzyl	31		495.34	466.11
121	3-trifluoromethyl phenyl	1-naphthyl	95	356 nM	395.38	396.15
122	3-trifluoromethyl phenyl	2-naphthyl	25		395.38	396.14
123	3-trifluoromethyl phenyl		36		384.36	385.13
124	3-trifluoromethyl phenyl		31		385.35	386.13
125	3-trifluoromethyl phenyl		94	232 nM	396.37	397.15
127	3-chlorophenyl	4-fluorophenyl	60		329.76	330.1
128	3-chlorophenyl	Phenyl	50		311.77	312.1
129	3-chlorophenyl		90	450 nM	362.81	363.1
130	3-chlorophenyl	3,5-dichlorophenyl	7		380.66	382
131	3-chlorophenyl	2-(1-pyrrolyl) phenyl	29		376.84	337.1
132	3- methoxyphenyl	4-fluorophenyl	38		325.34	326.1
133	3- methoxyphenyl	Phenyl	39		307.35	308.1

134	3-methoxyphenyl		82		358.40	359.1
135	3-methoxyphenyl	3,5-dichlorophenyl	27		376.24	376.0
136	3-methoxyphenyl	2-(1-pyrrolyl) phenyl	42		372.43	373.1
137	3-fluorophenyl	4-fluorophenyl	44		313.30	314.0
138	3-fluorophenyl	Phenyl	42		295.31	296.1
139	3-fluorophenyl		79		346.36	347.1
140	3-fluorophenyl	3,5-dichlorophenyl	29		364.20	364.0
141	3-fluorophenyl	2-(1-pyrrolyl) phenyl	42		360.39	361.1
142	3-chlorophenyl	3-trifluoromethyl phenyl		1.28 μ M	379.77	380.0
143	2-chloro-5-trifluoromethyl phenyl	4-fluorophenyl	15		397.58	398.1
144	2-chloro-5-trifluoromethyl phenyl	Phenyl	20		379.77	380.1
145	2-chloro-5-trifluoromethyl phenyl			324 nM	430.81	431.4
146	2-chloro-5-trifluoromethyl phenyl	3,5-dichlorophenyl	30		448.66	449.9
147	2-chloro-5-trifluoromethyl phenyl	2-(1-pyrrolyl) phenyl	21		444.84	445.1

148	2-chloro-5-trifluoromethyl phenyl		3		403.37	404.11
149	3-trifluoromethyl phenyl		35		435.45	434.3
150	3-trifluoromethyl phenyl		26		465.43	464.0
151	3-trifluoromethyl phenyl		18		477.44	477.44
152	3-trifluoromethyl phenyl			367 nM	410.40	411.6
153	3-trifluoromethyl phenyl		25		400.40	401.4
154	3-trifluoromethyl phenyl	3-nitrophenyl		1 μ M	390.32	391.11
155	3-trifluoromethyl phenyl		23		584.66	585.0
156	3-trifluoromethyl phenyl	3-aminophenyl	53		360.34	361.13
157	3-trifluoromethyl phenyl	3-benzoylphenyl		2 μ M	449.43	450.1
158	3-trifluoromethyl phenyl		52		530.52	531.3

EXAMPLE 4**IN VIVO ASSAY****Rodent Feeding Model:****5 Measurement of Food Intake in Food-Deprived Rats**

Male Long-Evans rats (180-200 grams) were housed individually and were maintained on a once-a-day feeding schedule (i.e. 10 a.m. until 4 p.m.) for five days following quarantine to allow the animals to acclimate to feeding on powdered chow (#5002 PMI Certified Rodent Meal) during the allotted time. The chow was made available in an open jar, anchored in the cage by a wire, with a metal follower covering the food to minimize spillage. Water was available ad-libitum.

15 Animals were fasted for 18 hours prior to testing. At the end of the fasting period, animals were administered either compounds of the invention or vehicle. Vehicle and test compounds were administered either orally (5 mL/kg) 60 minutes prior to the experiment, or 30 minutes prior when given subcutaneously (1 mL/kg) or intraperitoneally (1 mL/kg). Compounds of the invention were administered orally as a suspension in aqueous 0.5% methylcellulose-0.4% Tween 80, or intraperitoneally as a solution or suspension in PEG 200; compound concentrations typically range from 1 mg/kg to 100 mg/kg, preferably from 10-30 mg/kg. Food intake was measured at 2, 4, and 6 hours after administration by weighing the special jar containing the food before the experiment and at the specified times. Upon completion of the experiment, all animals were given a one-week washout period before retesting.

Percent reduction of food consumption was calculated subtracting the grams of food consumed by the treated group from the grams of food consumed by the control group divided by the grams of food consumed by the control group, multiplied by 100.

$$\% \text{ change} = \frac{\text{Treatment} - \text{Vehicle}}{\text{Vehicle}} \times 100$$


- 5 A negative value indicated a reduction in food consumption and a positive value indicated an increase in food consumption.

Compound	Dose (mg/kg) (# rats)	<u>Food Consumption (grams)</u>		
		0-2 hrs (% change)	0-6 hrs (% change)	2-6 hrs (% change)
Vehicle PEG-2000	N=16	8.44 g	18.0 g	9.56 g
# 6	30 (i.p.)	4.5 g	12.10 g	7.63 g
	N=16	(-46.7%)	(-32.8%)	(-20.2%)
Vehicle PEG-2000	N=8	10.38 g	23.38	13.0 g
# 110	30 (i.p.)	6.88 g	14.25 g	7.38 g
	N=8	(-33.73%)	(-39.04%)	(-43.3%)

- 10 While the foregoing specification teaches the principles of the present invention, with examples provided for the purpose of illustration, it will be understood that the practice of the invention encompasses all of the usual variations, adaptations and/or modifications as come within the scope of the following claims and their equivalents.

What is Claimed:

1. A compound of the formula (I):



(I)

wherein

5

- R¹ is selected from the group consisting of C₂-C₈ alkyl, aryl, substituted aryl, arC₁-C₈ alkyl, substituted arC₁-C₈ alkyl, heteroaryl, substituted heteroaryl, C₃-C₈ cycloalkyl, heteroC₃-C₈cycloalkyl, fluorinated C₁-C₈ alkyl, cyanoC₁-C₈alkyl and hydroxyC₁-C₈alkyl; wherein the aryl, aralkyl or heteroaryl group is substituted with one or more substituents independently selected from halogen, C₁-C₈ alkyl, C₁-C₈ alkoxy, fluorinated C₁-C₈ alkyl, fluorinated C₁-C₈ alkoxy, nitro, amino, amido, N-C₁-C₈alkylamido, N,N-di(C₁-C₈alkyl)amido, C₁-C₈ alkylsulfonyl, sulfonamido, N-C₁-C₈alkylsulfonamido, N,N-di(C₁-C₈alkyl)sulfonamido, C₁-C₈alkylsulfonylamino, or
- 10
- 15 C₁-C₈alkylcarbonylamino;

- R² is selected from the group consisting of di(C₁-C₆ alkyl)amino-C₁-C₆ alkyl, unsubstituted or substituted heteroarylC₀-C₆ alkyl, unsubstituted or substituted heterocycloalkylC₀-C₆ alkyl, unsubstituted or substituted aryl, unsubstituted or substituted arC₁-C₆ alkyl, and unsubstituted or substituted C₃-C₈ cycloalkylC₁-C₆ alkyl where the substituent is arylsulfonamidoC₁-C₆ alkyl;
- 20

wherein

- (e) the substituents on the aryl group are one or more substituents independently selected from the group consisting of halogen, nitro, amino, substituted amino where the substituents on
- 25

the amino are one or two of C₁-C₈ alkyl or phenyl; fluorinated C₁-C₆ alkyl, fluorinated C₁-C₆ alkoxy, C₁-C₈ alkyl, substituted C₁-C₈ alkyl where the substituent on the alkyl is N-(C₁-C₆ alkyl)-N-(C₃-C₈ cycloalkyl)amino; C₁-C₈ alkoxy, arC₁-C₈ alkyl, arC₁-C₈ alkoxy, arylcarbonyl, phenyl, aminophenyl, C₁-C₆ alkylthio, substituted arylsulfonamido where the substituent on the arylsulfonamido is C₁-C₆ alkyl or C₁-C₆ alkoxy; heteroaryl, substituted heteroaryl where the substituent on the heteroaryl is C₁-C₆ alkyl, heterocycloalkyl and oxo;

(f) the substituents on the aralkyl group are one or more substituents independently selected from halogen or fluorinated C₁-C₄ alkyl;

(g) the substituents on the heterocycloalkyl group are one or more substituents independently selected from C₁-C₆ alkyl or arC₁-C₈ alkyl;

(h) the substituents on the heteroaryl group are one or more substituents independently selected from oxo, C₁-C₆ alkyl or halogen;

R³ is selected from the group consisting of hydrogen, halogen, C₁-C₈ alkyl, arC₁-C₈ alkyl, heteroarylC₁-C₈ alkyl, C₁-C₈ alkoxy, fluorinated C₁-C₈ alkyl, fluorinated C₁-C₈ alkoxy, aminoC₁-C₈ alkyl and C₁-C₈ alkylaminoC₁-C₈ alkyl;

R⁴ is selected from the group consisting of halogen, C₁-C₈ alkyl, arC₁-C₈ alkyl, heteroarylC₁-C₈ alkyl, fluorinated C₁-C₈ alkyl, aminoC₁-C₈ alkyl and C₁-C₈ alkylaminoC₁-C₈ alkyl;

R⁵ is selected from the group consisting of hydrogen and C₁-C₈ alkyl;

provided that when R¹ is phenyl or benzyl, and R³ is hydrogen, and R⁴ is

5 methyl, and R⁵ is hydrogen, then R² is selected from di(C₁-C₆ alkyl)aminoC₁-C₆ alkyl, unsubstituted or substituted heteroarylC₀-C₆ alkyl, unsubstituted or substituted heterocycloalkylC₀-C₆ alkyl, substituted arC₁-C₆ alkyl, unsubstituted or substituted C₃-C₈ cycloalkylC₁-C₆ alkyl where the substituent is
10 arylsulfonamidoC₁-C₆ alkyl, and substituted aryl wherein the substituents on the aryl group are one or more substituents independently selected from the group consisting of halogen, nitro, amino, substituted amino where the substituents on the amino are one or two of C₁-C₆ alkyl or phenyl; fluorinated C₁-C₆ alkyl, fluorinated C₁-C₆ alkoxy, C₁-C₆ alkyl (provided that R² is not di(C₁-C₄ alkyl)phenyl), substituted C₁-C₆ alkyl where the substituent on the alkyl is N-(C₁-C₆ alkyl)-N-(C₃-C₈ cycloalkyl)amino; C₁-C₈ alkoxy, arC₁-C₈ alkyl, arC₁-C₈ alkoxy, phenyl, aminophenyl, C₁-C₆ alkylthio, substituted arylsulfonamido where the substituent on the arylsulfonamido is C₁-C₆ alkyl or C₁-C₆ alkoxy; heteroaryl, substituted heteroaryl where the substituent on the heteroaryl is C₁-
15 C₆ alkyl; heterocycloalkyl and oxo;

and pharmaceutically acceptable salts thereof.

2. The compound of Claim 1, wherein R⁵ is hydrogen.
- 20 3. The compound of Claim 1, wherein

R³ is hydrogen, R⁴ is C₁-C₄ alkyl, and R⁵ is hydrogen or C₁-C₄ alkyl;

- 25 provided that when R¹ is phenyl or benzyl, and R⁴ is methyl, then R² is selected from di(C₁-C₆ alkyl)aminoC₁-C₆ alkyl, unsubstituted or substituted heteroarylC₀-C₆ alkyl, unsubstituted or substituted heterocycloalkylC₀-C₆ alkyl, substituted arC₁-C₆ alkyl, unsubstituted or substituted C₃-C₈ cycloalkylC₁-C₆ alkyl where the substituent is arylsulfonamidoC₁-C₆ alkyl, and substituted aryl
30 wherein the substituents on the aryl group are one or more substituents independently selected from the group consisting of halogen, nitro, amino, substituted amino where the substituents on the amino are one or two of C₁-C₆

alkyl or phenyl; fluorinated C₁-C₆ alkyl, fluorinated C₁-C₆ alkoxy, C₁-C₆ alkyl (provided that R² is not dimethylphenyl), substituted C₁-C₆ alkyl where the substituent on the alkyl is N-(C₁-C₆ alkyl)-N-(C₃-C₈ cycloalkyl)amino; C₁-C₈ alkoxy, arC₁-C₈ alkyl, arC₁-C₈ alkoxy, phenyl, aminophenyl, C₁-C₆ alkylthio, substituted arylsulfonamido where the substituent on the arylsulfonamido is C₁-C₆ alkyl or C₁-C₆ alkoxy; heteroaryl, substituted heteroaryl where the substituent on the heteroaryl is C₁-C₆ alkyl; heterocycloalkyl and oxo;

and pharmaceutically acceptable salts thereof.

10

4. The compound of Claim 2, wherein

R¹ is selected from the group consisting of pyridyl and substituted phenyl, wherein the phenyl is substituted with one to three substituents independently selected from C₁-C₄ alkyl, C₁-C₄ alkoxy, fluorinated C₁-C₄ alkyl, fluorinated C₁-C₄ alkoxy or halogen;

R² is selected from the group consisting of unsubstituted phenyl or substituted phenyl, where the substituents on the phenyl are independently selected from one to three of halogen, C₁-C₄ alkoxy, pyrrolyl, morpholino, aminophenyl, benzimidazolyl, fluorinated C₁-C₄ alkyl, fluorinated C₁-C₄ alkoxy, C₁-C₄ alkylthio, nitro, C₁-C₄ alkyl, amino, C₁-C₄ alkylamino, phenyl amino, piperidinyl, phenyl, benzyl, benzoyl, benzyloxy, C₁-C₄ alkyl substituted benthiazolyl, C₁-C₄ alkyl substituted phenylsulfonamido, C₁-C₄ alkoxy substituted phenylsulfonamido or N-(C₁-C₄ alkyl)-N-(C₃-C₈ cycloalkyl)amino C₁-C₄ alkyl; unsubstituted or substituted benzyl, where the substituents on the benzyl are independently selected from one to three of halogen or fluorinated C₁-C₄ alkyl; naphthyl, 3,4-methylenedioxyphenyl, 3,4-ethylenedioxyphenyl, 5,6,7,8-tetrahydronaphthyl, indanyl, pyridyl, pyrazinyl, purinyl, anthraquinoliny, fluoro- substituted benzthiazolyl, indolyl, indolylC₁-C₄ alkyl, quinoliny, fluorenyl, indazolyl, 2,1,3-benzthiasoyl, isoquinoliny, methyl substituted isoquinoliny, N-benzyl-4-piperidinyl, methyl substituted piperizinyl, 3-(2-methyl piperidinyl)

propyl, di(C₁-C₄ alkyl)aminoC₁-C₄ alkyl, naphthylsulfonamidoC₁-C₄ alkyl substituted cyclohexylC₁-C₄ alkyl, 5-(3,4-benzocoumarinyl) and 5-(1,2,3,4-tetrahydroisoquinolyl);

- 5 R³ is selected from the group consisting of hydrogen, halogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, fluorinated C₁-C₄ alkyl and fluorinated C₁-C₄ alkoxy;

 R⁴ is selected from the group consisting of hydrogen, C₁-C₄ alkyl and fluorinated C₁-C₄ alkyl;

10

and pharmaceutically acceptable salts thereof.

5. The compound of Claim 3, wherein,

- 15 R¹ is selected from the group consisting of pyridyl and substituted phenyl, wherein the phenyl is substituted with one to three substituents independently selected from C₁-C₄ alkyl, C₁-C₄ alkoxy, fluorinated C₁-C₄ alkyl, fluorinated C₁-C₄ alkoxy or halogen;

- 20 R² is selected from the group consisting of unsubstituted phenyl or substituted phenyl, where the substituents on the phenyl are independently selected from one to three of halogen, C₁-C₄ alkoxy, pyrrolyl, morpholino, aminophenyl, benzimidazolyl, fluorinated C₁-C₄ alkyl, fluorinated C₁-C₄ alkoxy, C₁-C₄ alkylthio, nitro, C₁-C₄ alkyl, amino, C₁-C₄ alkylamino, phenyl amino, piperidinyl, phenyl, benzyl, benzoyl, benzyloxy, C₁-C₄ alkyl substituted benthiazolyl, C₁-C₄ alkyl substituted phenylsulfonamido, C₁-C₄ alkoxy substituted phenylsulfonamido or N-(C₁-C₄ alkyl)-N-(C₃-C₆ cycloalkyl)amino C₁-C₄ alkyl; unsubstituted or substituted benzyl, where the substituents on the benzyl are independently selected from one to three of halogen or fluorinated C₁-C₄ alkyl; naphthyl, 3,4-methylenedioxyphenyl, 3,4-ethylenedioxyphenyl, 5,6,7,8-tetrahydronaphthyl, indanyl, pyridyl, pyrazinyl, purinyl, anthraquinoliny, fluoro- substituted benzthiazolyl, indolyl, indolylC₁-C₄ alkyl, quinoliny, fluorenyl,
- 25
- 30

indazolyl, 2,1,3-benzthiasoyl, isoquinolinyl, methyl substituted isoquinolinyl, N-benzyl-4-piperidinyl, methyl substituted piperiziny, 3-(2-methyl piperidinyl) propyl, di(C₁-C₄ alkyl)aminoC₁-C₄ alkyl, naphthylsulfonamidoC₁-C₄ alkyl substituted cyclohexylC₁-C₄ alkyl, 5-(3,4-benzocoumarinyl) and 5-(1,2,3,4-tetrahydroisoquinolyl);

R⁵ is hydrogen;

and pharmaceutically acceptable salts thereof.

10

6. The compound of Claim 5 wherein

R¹ is selected from the group consisting of phenyl, 3-tolyl, 3-trifluoromethylphenyl, 3,5-dichlorophenyl, 3-chlorophenyl, and 2-chloro-5-trifluoromethylphenyl;

15

R² is selected from the group consisting of 4-fluorophenyl, 3,5-dichlorophenyl, 2,3-dichlorophenyl, 2-(1-pyrrolyl) phenyl, 2-(aminophenyl)phenyl, 3-trifluoromethoxyphenyl, 2-methylthio, 3-tolyl, 2-tolyl, 2,6-difluorophenyl, 2,3,4-trifluorophenyl, 2,3,6-trifluorophenyl, 3-chlorophenyl, 3-bromophenyl, 3-iodophenyl, 3-chloro-5-fluorophenyl, 3-methoxy-5-trifluoromethylphenyl, 3-trifluoromethyl-4-fluorophenyl, 5,6,7,8-tetrahydronaphthyl, 2-benzylphenyl, phenyl, 3-trifluoromethylphenyl, 5-isoquinolinyl, 2-(N-methylamino)phenyl, 1-naphthyl, 5-quinolinyl, 5-(3-methyl)isoquinolinyl and 3-nitrophenyl;

25

R⁴ is methyl;

and pharmaceutically acceptable salts thereof.

30

7. The compound of Claim 6, wherein

R¹ is selected from the group consisting of 3-trifluoromethylphenyl, 3,5-dichlorophenyl and 3-tolyl;

5 R² is selected from the group consisting of 3,5-dichlorophenyl, 2-(aminophenyl)-phenyl, 2,6-difluorophenyl, 2,3,6-trifluorophenyl, 5,6,7,8-tetrahydronaphthyl, 2-benzylphenyl, 3-trifluoromethylphenyl, 5-isoquinolyl and 5-quinolinyl;

and pharmaceutically acceptable salts thereof.

10

8. The compound of Claim 7, wherein R¹ is 3-trifluoromethylphenyl and R² is selected from 3,5-dichlorophenyl or 5-isoquinolyl and pharmaceutically acceptable salts thereof.

15

9. A pharmaceutical composition comprising a compound of Claim 1 and a pharmaceutically acceptable carrier.

10. A pharmaceutical composition made by mixing a compound of Claim 1 and a pharmaceutically acceptable carrier.

20

11. A process for making a pharmaceutical composition comprising mixing a compound of Claim 1 and a pharmaceutically acceptable carrier.

25

12. A method of treating a disorder mediated by the NPY Y5 receptor comprising administering to a subject in need of such treatment a therapeutically effective amount of a compound of claim 1.

13. The method of Claim 12, wherein the therapeutically effective amount is between about 0.03 and about 300 mg/kg per day.

30

14. The method of Claim 13, wherein the disorder is selected from an eating disorder, obesity, bulimia nervosa, diabetes, binge eating, anorexia

nervosa, dyslipidimia, hypertension, memory loss, epileptic seizures, migraine, sleep disturbances, pain, sexual/reproductive disorders, depression, anxiety, cerebral hemorrhage, shock, congestive heart failure, nasal congestion or diarrhea.

5

15. A method of treating a disorder selected from an eating disorder, obesity, bulimia nervosa, diabetes, binge eating, anorexia nervosa, dyslipidimia, hypertension, memory loss, epileptic seizures, migraine, sleep disturbances, pain, sexual/reproductive disorders, depression, anxiety, cerebral hemorrhage, shock, congestive heart failure, nasal congestion or diarrhea in a subject in need thereof, comprising administering to the subject a therapeutically effective amount of a compound of Claim 1.

10

- 15 16. The method of claim 15, wherein the therapeutically effective amount is between about 0.03 and about 300 mg/kg per day.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 00/11903

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D401/12 C07D231/14 C07D403/12 C07D405/12 C07D473/34
C07D417/12 A61K31/415 A61K31/4155

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

CHEM ABS Data, EPO-Internal, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 98 57951 A (DU PONT MERCK PHARMA) 23 December 1998 (1998-12-23) claims; example 80; table 2 ---	1,2,9-11
X	DE 16 70 945 A (BAYER) 2 March 1972 (1972-03-02) example 27 ---	1,2,9-11
X	LAN, RUOXI ET AL: "Structure-Activity Relationships of Pyrazole Derivatives as Cannabinoid Receptor Antagonists" J. MED. CHEM. (1999), 42(4), 769-776 , XP002145465 compound 16 ---	1,2,9-11
X	EP 0 289 879 A (MITSUBISHI CHEM IND) 9 November 1988 (1988-11-09) claim 1; examples ---	1,2
	-/-	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

23 August 2000

Date of mailing of the international search report

19/09/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3018

Authorized officer

De Jong, B

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 00/11903

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 98 25907 A (BANYU PHARMA CO LTD ;FUKURODA TAKAHIRO (JP); KANATANI AKIO (JP); F) 18 June 1998 (1998-06-18) cited in the application abstract -& EP 0 945 438 A 29 September 1999 (1999-09-29)	1,9-16
A	WO 98 25908 A (BANYU PHARMA CO LTD ;FUKURODA TAKAHIRO (JP); KANATANI AKIO (JP); F) 18 June 1998 (1998-06-18) cited in the application abstract -& EP 0 945 439 A 29 September 1999 (1999-09-29)	1,9-16
A	WO 98 27063 A (BANYU PHARMA CO LTD ;FUKURODA TAKAHIRO (JP); KANATANI AKIO (JP); F) 25 June 1998 (1998-06-25) cited in the application abstract -& EP 0 945 440 A 29 September 1999 (1999-09-29)	1,9-16

INTERNATIONAL SEARCH REPORT

Information on patent family members

Int. l. Application No

PCT/US 00/11903

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9857951 A	23-12-1998	AU 7976898 A EP 0991638 A HR 980333 A LT 99147 A NO 995965 A	04-01-1999 12-04-2000 28-02-1999 25-04-2000 03-12-1999
DE 1670945 A	02-03-1972	AT 288421 B AT 287733 B BE 723336 A CA 932728 A CH 513873 A CH 516578 A CH 516579 A CH 516580 A CS 158230 B CS 158229 B ES 359830 A ES 363815 A FI 50414 B FR 8220 M FR 1601129 A GB 1176288 A IE 32534 B NL 6815633 A NO 128111 B SE 362249 B SE 362430 B SE 365220 B US 3718660 A YU 253868 A, B SE 336338 B	15-01-1971 10-02-1971 05-05-1969 28-08-1973 15-10-1971 15-12-1971 15-12-1971 15-12-1971 15-10-1974 15-10-1974 16-06-1970 16-12-1970 01-12-1975 21-09-1970 10-08-1970 01-01-1970 05-09-1973 06-05-1969 01-10-1973 03-12-1973 10-12-1973 18-03-1974 27-02-1973 31-03-1976 05-07-1971
EP 0289879 A	09-11-1988	AT 86979 T BR 8802011 A CA 1330342 A CN 1028713 B DE 3879262 A DE 3879262 T ES 2053611 T IN 166898 A JP 1025763 A JP 1838897 C JP 5051582 B KR 9509363 B PT 87302 A, B US 4950668 A ZA 8802738 A	15-04-1993 21-11-1989 21-06-1994 07-06-1995 22-04-1993 02-09-1993 01-08-1994 04-08-1990 27-01-1989 25-04-1994 03-08-1993 21-08-1995 01-05-1988 21-08-1990 17-10-1988
WO 9825907 A	18-06-1998	AU 5410498 A EP 0945438 A US 6057335 A	03-07-1998 29-09-1999 02-05-2000
WO 9825908 A	18-06-1998	AU 5410598 A EP 0945439 A	03-07-1998 29-09-1999
WO 9827063 A	25-06-1998	AU 7738198 A EP 0945440 A	15-07-1998 29-09-1999